

## Hemoglobin S-nitrosylation plays an essential role in cardioprotection

Rongli Zhang, ... , James D. Reynolds, Jonathan S. Stamler

*J Clin Invest.* 2016;126(12):4654-4658. <https://doi.org/10.1172/JCI90425>.

Brief Report

Cardiology

Vascular biology

Homeostatic control of tissue oxygenation is achieved largely through changes in blood flow that are regulated by the classic physiological response of hypoxic vasodilation. The role of nitric oxide (NO) in the control of blood flow is a central tenet of cardiovascular biology. However, extensive evidence now indicates that hypoxic vasodilation entails S-nitrosothiol-based (SNO-based) vasoactivity (rather than NO per se) and that this activity is conveyed substantially by the  $\beta$ Cys93 residue in hemoglobin. Thus, tissue oxygenation in the respiratory cycle is dependent on S-nitrosohemoglobin. This perspective predicts that red blood cells (RBCs) may play an important but previously undescribed role in cardioprotection. Here, we have found that cardiac injury and mortality in models of myocardial infarction and heart failure were greatly enhanced in mice lacking  $\beta$ Cys93 S-nitrosylation. In addition,  $\beta$ Cys93 mutant mice exhibited adaptive collateralization of cardiac vasculature that mitigated ischemic injury and predicted outcomes after myocardial infarction. Enhanced myopathic injury and mortality across different etiologies in the absence of  $\beta$ Cys93 confirm the central cardiovascular role of RBC-derived SNO-based vasoactivity and point to a potential locus of therapeutic intervention. Our findings also suggest the possibility that RBCs may play a previously unappreciated role in heart disease.

Find the latest version:

<https://jci.me/90425/pdf>



# Hemoglobin S-nitrosylation plays an essential role in cardioprotection

Rongli Zhang,<sup>1,2</sup> Douglas T. Hess,<sup>1,2</sup> James D. Reynolds,<sup>1,3,4</sup> and Jonathan S. Stamler<sup>1,2,4</sup>

<sup>1</sup>Institute for Transformative Molecular Medicine, Case Western Reserve University School of Medicine and University Hospitals Cleveland Medical Center, Cleveland, Ohio, USA. <sup>2</sup>Department of Medicine, Case Western Reserve University School of Medicine, Cleveland, Ohio, USA. <sup>3</sup>Department of Anesthesiology and Perioperative Medicine, Case Western Reserve University School of Medicine, Cleveland, Ohio, USA. <sup>4</sup>Harrington Discovery Institute, University Hospitals Cleveland Medical Center, Cleveland, Ohio, USA.

**Homeostatic control of tissue oxygenation is achieved largely through changes in blood flow that are regulated by the classic physiological response of hypoxic vasodilation. The role of nitric oxide (NO) in the control of blood flow is a central tenet of cardiovascular biology. However, extensive evidence now indicates that hypoxic vasodilation entails S-nitrosothiol-based (SNO-based) vasoactivity (rather than NO per se) and that this activity is conveyed substantially by the  $\beta$ Cys93 residue in hemoglobin. Thus, tissue oxygenation in the respiratory cycle is dependent on S-nitrosohemoglobin. This perspective predicts that red blood cells (RBCs) may play an important but previously undescribed role in cardioprotection. Here, we have found that cardiac injury and mortality in models of myocardial infarction and heart failure were greatly enhanced in mice lacking  $\beta$ Cys93 S-nitrosylation. In addition,  $\beta$ Cys93 mutant mice exhibited adaptive collateralization of cardiac vasculature that mitigated ischemic injury and predicted outcomes after myocardial infarction. Enhanced myopathic injury and mortality across different etiologies in the absence of  $\beta$ Cys93 confirm the central cardiovascular role of RBC-derived SNO-based vasoactivity and point to a potential locus of therapeutic intervention. Our findings also suggest the possibility that RBCs may play a previously unappreciated role in heart disease.**

## Introduction

Control of blood flow in the microvasculature (i.e., tissue perfusion) in response to altered O<sub>2</sub> consumption is governed by an autoregulatory mechanism whereby flow is coupled to tissue pO<sub>2</sub> (1). It has been shown that red blood cells (RBCs) can themselves provide the signal that couples flow to pO<sub>2</sub>, and it was proposed that this signal is mediated by the pO<sub>2</sub>-regulated release of nitric oxide-derived (NO-derived) vasodilatory activity that is conveyed by hemoglobin (Hb) S-nitrosylated at Cys93 of the  $\beta$ -chain (SNO-Hb) (2, 3) (reviewed in ref. 4). This proposal was recently confirmed (5) by analyses of engineered mice in which the critical Cys that mediates vasodilation by Hb was replaced by Ala ( $\beta$ Cys93Ala) (6). In the absence of  $\beta$ Cys93, peripheral blood flow and tissue oxygenation are markedly decreased at normoxia and decline excessively during hypoxia (5). In addition, the  $\beta$ Cys93Ala mutation results in myocardial ischemia under basal, normoxic conditions and in acute cardiac decompensation and enhanced mortality during transient hypoxia (5). Therefore, the ability of mammals to oxygenate tissues, including oxygenation of the heart required for normal myocardial function, is evidently dependent in large part upon SNO-Hb. Tissue oxygenation is thus governed not only by the O<sub>2</sub> content, but also by the SNO

content of Hb, consistent with the strict evolutionary conservation of  $\beta$ Cys93 in all mammals (7).


The most common causes of morbidity and mortality in Western cultures are cardiovascular in origin. Both extent of injury and outcomes in the most prevalent cardiac diseases, i.e., acute myocardial infarction (MI) and heart failure, reflect uncoupling of blood flow from metabolic demand (8, 9). Our recent findings in  $\beta$ Cys93Ala animals (5) support a new perspective in which RBCs subserve blood-flow regulation previously ascribed to the endothelium. To test the resulting prediction that regulation of blood flow by RBCs would play a role in response to cardiac insult, we extended our analysis of  $\beta$ Cys93Ala animals to well-established experimental murine models of MI, induced by ischemia-reperfusion (IR) injury (10), and pressure overload-induced cardiomyopathy, induced by transaortic constriction (TAC) (11). Our results demonstrate that  $\beta$ Cys93 in Hb plays an essential role in the outcome of both insults.

## Results and Discussion


The mice employed comprised 3 genetically engineered strains expressing “humanized” RBC Hb (6), which we designated  $\gamma$  $\beta$ C93 (control mice),  $\gamma$  $\beta$ C93A, and  $\beta$ C93A. In all engineered strains, the murine  $\alpha$  and  $\beta$  subunits of RBC Hb were replaced with the human versions. In  $\gamma$  $\beta$ C93 and  $\gamma$  $\beta$ C93A mice (genetically matched with the exception of  $\beta$ Cys93), RBCs also contained the human  $\gamma$  (fetal) subunit; in all 3 strains, RBCs retained the murine  $\gamma$  subunit, but  $\gamma$  subunits remained minor species (5, 12).  $\beta$ Cys93 was replaced with Ala in  $\beta$ C93A and  $\gamma$  $\beta$ C93A mice. It has previously been shown that total Hb concentration, blood O<sub>2</sub> content, O<sub>2</sub>-binding curves, blood pressure, and heart rate do not differ significantly, or meaningfully,

### ► Related Commentary: p. 4402

**Conflict of interest:** J.S. Stamler has financial interests in Nivalis Therapeutics Inc., Adamas Pharmaceuticals, and LifeHealth. J.D. Reynolds has a financial interest in Miach Medical Innovations. J.S. Stamler and J.D. Reynolds have licensed technology to Novartis.  
**Submitted:** August 30, 2016; **Accepted:** October 6, 2016.  
**Reference information:** *J Clin Invest.* 2016;126(12):4654–4658. doi:10.1172/JCI90425.



**Figure 1. Enhanced MI-induced cardiac injury and mortality in the absence of  $\beta$ Cys93.** (A) IR-induced mortality is greatly enhanced in mice lacking  $\beta$ Cys93 (C93A =  $\gamma\beta$ C93A and  $\beta$ C93A) versus control mice (C93 =  $\gamma\beta$ C93). In panels A, C, and D, numbers within histograms specify absolute ratios for each observation (B) Representative angiography illustrating collateralized circulation of the LV free wall to supply the apex in a mutant  $\beta$ C93A mouse. Lower (left;  $\times 6.7$ ) and higher (right;  $\times 15.8$ ) magnification views are shown. (C) Collateralization supplying the LV is significantly more likely in  $\gamma\beta$ C93A and  $\beta$ C93A versus  $\gamma\beta$ C93 mice.  $*P < 0.0001$  for  $\gamma\beta$ C93 versus  $\gamma\beta$ C93A and  $**P = 0.0043$  for  $\gamma\beta$ C93 versus  $\beta$ C93A, respectively, by Fisher's exact test (2-sided). The difference in collateralization between  $\gamma\beta$ C93A and  $\beta$ C93A mice was not significant ( $P > 0.05$ ). (D) Mortality during 24 hours of IR is determined by the presence versus absence of collateral circulatory supply of the LV and is enhanced in  $\gamma\beta$ C93A and  $\beta$ C93A versus  $\gamma\beta$ C93 mice.  $*P < 0.0001$  vs. collateral by Fisher's exact test (2-sided). (E) Representative Evans blue/TTC-stained LV slices illustrating areas of recovery (red) and necrosis (white) in a  $\gamma\beta$ C93 mouse (left) and in a  $\gamma\beta$ C93A mouse (right). Scale bar: 2 mm. (F) In  $\gamma\beta$ C93A versus  $\gamma\beta$ C93 control mice surviving 24 hours after IR, the area of recovery (Rec/LV; LV, total LV area) is significantly diminished and the area of necrosis (AON/LV) and the ratio of area of necrosis/area at risk (AON/AAR) are significantly enhanced, and enhanced injury was evident in the absence of  $\beta$ C93 even when collateralization was present. (G) Echocardiography of surviving mice reveals that LV ejection fraction (EF) and fractional shortening (FS) are significantly diminished in  $\gamma\beta$ C93A versus  $\gamma\beta$ C93 control mice, and these differences were evident even when collateralization was present. (H) Thickening of the LV anterior and posterior walls in the absence of collateral blood supply is significantly diminished in  $\gamma\beta$ C93A versus  $\gamma\beta$ C93 mice. In F–H,  $n = 3$ –7 for mice with collateral blood supply and  $n = 3$ –9 for mice without collateral blood supply.  $*P < 0.05$  vs.  $\gamma\beta$ C93 by Student's  $t$  test (2-tailed).



**Figure 2. Assessment of acute effects of pressure overload (TAC) on cardiac function in C93A mutant animals.**  $\gamma\beta C93$  and  $\gamma\beta C93A$  mice were compared at 2 days. (A–C) Echocardiography revealed significantly decreased ejection fraction (A), fractional shortening (B), and cardiac output (C) in  $\gamma\beta C93A$  versus  $\gamma\beta C93$  mice. (D) End systolic diameter (end systolic diam.) was significantly greater in  $\gamma\beta C93A$  versus  $\gamma\beta C93$  mice. (E and F) The ratios of heart weight with respect to body weight (HW/BW) (E) and lung weight with respect to body weight (LW/BW) (F) were significantly increased in  $\gamma\beta C93A$  versus  $\gamma\beta C93$  mice.  $n = 5-7$ . \* $P < 0.05$  vs.  $\gamma\beta C93$  by Student's  $t$  test (2-tailed).

among these strains under basal conditions (5, 6, 12). We focused our analysis on comparisons between the  $\gamma\beta C93$  and  $\gamma\beta C93A$  strains, isogenic except for the mutation of  $\beta C93$ . However, we have shown that the cardiovascular phenotypes of  $\gamma\beta C93$  or  $\beta C93A$  (versus  $\gamma\beta C93$ ) mice are similar (5), and because of difficulties with breeding and therefore maintaining colonies of sufficient numbers for studies (5), we included comparisons with  $\beta C93A$  mice where those comparisons help clarify interpretations of results.

MI was induced by occlusion of the left coronary artery (LCA) for 30 minutes, followed by reperfusion for 24 hours (IR). All mice recovered from the surgery, but within 24 hours of IR, approximately 14% (2/14) of control mice died, whereas approximately 32% (12/38) of mice lacking  $\beta C93$  died (Figure 1A). Thus, delivery of vasodilatory activity via  $\beta C93$  is cardioprotective in the setting of IR-induced MI.

Employing angiography or Evans blue staining (see Supplemental Methods; supplemental material available online with this article; doi:10.1172/JCI90425DS1), we observed that major collaterals of the LCA or of the right coronary artery (RCA) traversed the left ventricle (LV) to terminate in the apex (Figure 1B) in 14% (5/36) of  $\gamma\beta C93$  control mice (Figure 1C) and that collateralization was greatly enhanced in  $\gamma\beta C93A$  (63%; 19/30) and  $\beta C93A$  (44%; 22/50) mutant mice (Figure 1C). The incidence of collateralization did not differ significantly between  $\gamma\beta C93A$  and  $\beta C93A$  mice (Figure 1C). Collateralization of the LV from the LCA versus the RCA, which did not occur together, was equally likely in all strains. Enhanced collateralization in  $\gamma\beta C93A$  and  $\beta C93A$  mice was consistent with compensation for inadequate myocardial oxygenation in the absence of  $\beta C93$ , previously demonstrated in  $\beta C93A$  and  $\gamma\beta C93A$  mice (5).

We examined the effects of collateralization on IR-induced mortality. As shown in Figure 1D, no mice died during the reperfusion interval when collateralization was present ( $\gamma\beta C93$ ,  $n = 3$ ;  $\gamma\beta C93A$ ,  $n = 7$ ;  $\beta C93A$ ,  $n = 11$ ). However, in the absence of collateralization, whereas only 18% of  $\gamma\beta C93$  control mice died (2/11), mortality was greatly enhanced in  $\gamma\beta C93A$  (57%; 4/7) and  $\beta C93A$  (62%; 8/13) mutant mice. Mortality in the absence of collateralization did not differ significantly between  $\gamma\beta C93A$  and  $\beta C93A$  mutant mice. Thus, compensatory collateralization protects from IR-induced death even in the absence of  $\beta C93$ , but without collateralization, the absence of  $\beta C93$  results in greatly enhanced mortality.

We obtained standard measures of injury of the LV following IR in  $\gamma\beta C93$  and  $\gamma\beta C93A$  mice that survived the 24-hour reper-


fusion interval, employing staining with Evans blue and 2,3,5-triphenyltetrazolium chloride (TTC): area of recovery, area at risk, and area of necrosis (Figure 1E). In  $\gamma\beta C93A$  versus  $\gamma\beta C93$  mice in the absence of collateralization, the area of recovery was significantly diminished, and the area of necrosis and the ratio of the area of necrosis to the area at risk was significantly enlarged (whereas the area at risk was not significantly different); notably, enhanced injury was evident in the absence of  $\beta C93$  even when collateralization was present (Figure 1F).

We also assessed cardiac function by echocardiography in  $\gamma\beta C93$  and  $\gamma\beta C93A$  mice that survived the 24-hour reperfusion interval. We measured LV ejection fraction and fractional shortening (Figure 1G) as well as anterior and posterior LV wall thickening (Figure 1H). All of these measures declined in  $\gamma\beta C93A$  versus  $\gamma\beta C93$  mice (Figure 1, G and H), and significant declines in ejection fraction and fractional shortening were evident in the absence of  $\beta C93$  even when collateralization was present (Figure 1G). Thus, the absence of  $\beta C93$  induces remodeling of cardiac vasculature and heart and, absent vascular remodeling, greatly increases mortality associated with MI; cardiac injury in the absence of  $\beta C93$  is similarly increased.

We next asked whether protection by SNO-Hb extended to heart failure of nonischemic etiology and in particular to pressure overload-induced cardiomyopathy (11). Compensatory cardiac hypertrophy in response to increased workload is normally accompanied by angiogenesis to maintain oxygenation of hypertrophied cardiomyocytes, and inhibition of angiogenesis leads to heart failure (13). Therefore, it might be predicted that deficient oxygenation resulting from the absence of  $\beta C93$  would be deleterious.

Pressure overload-induced cardiac hypertrophy was induced by partial occlusion of the aorta (TAC). Two days after TAC, echocardiography revealed significantly greater impairment of LV function in  $\gamma\beta C93A$  versus  $\gamma\beta C93$  mice across all measures, including decreased ejection fraction (Figure 2A), fractional shortening (Figure 2B), and cardiac output (Figure 2C) as well as increased end systolic diameter (LV dilation) (Figure 2D). In addition, the ratios of heart weight (Figure 2E) and lung weight (Figure 2F) to body weight were elevated, indicative of cardiac hypertrophy and pulmonary edema, respectively.

Cohorts of  $\gamma\beta C93$ ,  $\gamma\beta C93A$ , and  $\beta C93A$  mice were followed for 30 days after TAC. Within 5 days after TAC, a single  $\gamma\beta C93$  control animal of 18 mice subjected to TAC died (1/18; 5.6% mortality), whereas 7 of 11  $\gamma\beta C93A$  mice died (63.6% mortality) before



**Figure 3. Assessment of chronic effects of pressure overload (TAC) on mortality and injury in C93A mutant animals.** Comparisons among  $\gamma\beta\text{C93}$ ,  $\beta\text{C93A}$ , and  $\gamma\beta\text{C93A}$  mice were made following 4 weeks of TAC. (A) A Kaplan-Meier curve illustrates that, within 5 days after TAC, only 1 of 18  $\gamma\beta\text{C93}$  control animals died (5.6% mortality), whereas 7 of 11  $\gamma\beta\text{C93A}$  mice died (63.6% mortality) before the population stabilized. Also, 12 of 23  $\beta\text{C93A}$  mice died (52.2% mortality), with deaths accumulating over the full TAC time course. (B) Cumulative mortality over 4 weeks was significantly greater for  $\beta\text{C93A}$  and  $\gamma\beta\text{C93A}$  versus  $\gamma\beta\text{C93}$  mice and did not differ significantly between  $\beta\text{C93A}$  and  $\gamma\beta\text{C93A}$  mice. \* $P = 0.0019$  vs.  $\gamma\beta\text{C93}$  for  $\gamma\beta\text{C93A}$  and \*\* $P = 0.0014$  vs.  $\gamma\beta\text{C93}$  for  $\beta\text{C93A}$  by Fisher's exact test (2-sided). (C–F) Echocardiography in mice surviving the 4-week TAC interval revealed greater impairment of LV function in  $\beta\text{C93A}$  versus  $\gamma\beta\text{C93}$  mice with respect to ejection fraction (C), fractional shortening (D), cardiac output (E), and end systolic diameter (LV dilation) (F). (G and H) Ratios of heart weight with respect to body weight (G) and lung weight with respect to body weight (H) were significantly increased in  $\beta\text{C93A}$  versus  $\gamma\beta\text{C93}$  mice, indicative of increased myocardial hypertrophy and pulmonary edema, respectively. In C–H,  $n = 11$  ( $\gamma\beta\text{C93}$ ) or 12 ( $\beta\text{C93A}$ );  $P < 0.01$  by Student's *t* test (2-tailed). (I) Post-mortem examination revealed lung edema and pleural effusion (yellow arrows; a representative  $\beta\text{C93A}$  mouse is shown).

the population stabilized (Figures 3, A and B). Similarly, 12 of 23  $\beta\text{C93A}$  mice died (52.2% mortality), with deaths accumulating over the full post-TAC time course (Figure 3, A and B). Cumulative mortality did not differ significantly between  $\gamma\beta\text{C93A}$  and  $\beta\text{C93A}$  mice (Figure 3B).

We assessed cardiac function by echocardiography in surviving mice. The small population of survivors precluded meaningful analysis of the  $\gamma\beta\text{C93A}$  strain. Echocardiography revealed significantly greater impairment of LV function in  $\beta\text{C93A}$  versus  $\gamma\beta\text{C93}$  mice across all measures including ejection fraction (Figure 3C), fractional shortening (Figure 3D), and cardiac output (Figure 3E) as well as significantly increased end systolic diameter (Figure 3F). In addition,  $\beta\text{C93A}$  mice versus  $\gamma\beta\text{C93}$  mice exhibited significantly increased ratios of heart weight to body weight (Figure 3G) and lung weight to body weight (Figure 3H), again indicative of cardiac hypertrophy and pulmonary edema, respectively. Post-mortem examination confirmed lung edema and pleural effusion in  $\beta\text{C93A}$  mutant mice (Figure 3I), consistent with clinical heart failure.

Our findings establish a role for Hb  $\beta\text{C93}$  in ameliorating the consequences of both ischemic and pressure overload-induced cardiac insults and thus demonstrate the central cardiovascular role of RBC-derived SNO-based vasoactivity (2, 3, 5) in the context of cardiac injury. These findings are consistent with the demonstration that  $\beta\text{Cys93A}$  mutation results in myocardial hypoxia

under basal, normoxic conditions (5) by impairing the allosteric mechanism in Hb through which hypoxia is coupled to compensatory increases in blood flow (2–4). Hypoxia-coupled delivery of SNO-based vasodilatory activity from  $\beta\text{C93}$  in vivo would enhance myocardial perfusion following ischemic injury (and thus the area of recovery is diminished and mortality is enhanced following IR in the absence of  $\beta\text{Cys93}$ ) and would support the compensatory myocardial hypertrophy and wall stress induced by TAC (and thus cardiac function is diminished and mortality is enhanced during TAC in the absence of  $\beta\text{Cys93}$ ).

Remarkably,  $\beta\text{Cys93}$  is the only residue in Hb that is conserved across mammals and birds, other than residues that coordinate the  $\text{O}_2$ -carrying heme moiety (7). Our results provide further evidence for the evolutionary selective pressure that underlies this conservation, including adaptive growth of LV collateral circulation to compensate for tissue ischemia. Because deficient delivery of SNO-based vasoactivity by Hb exacerbates cardiac injury across multiple etiologies, whereas increased RBC-derived NO bioactivity may ameliorate cardiac injury (14), our results suggest the possibility that assessment of SNO-Hb might provide a novel biomarker of cardiac disease. Our findings further suggest that enhancing delivery of SNO-based vasodilatory activity from Hb  $\beta\text{Cys93}$  may represent a potential therapeutic strategy for improving oxygenation in acute coronary syndromes and heart failure and more general-



ly across multiple pathophysiologies that are characterized by deficient tissue oxygenation.

## Methods

**MI model.** Male mice at 9 to 16 weeks of age were subjected to IR induced by transient occlusion (30 minutes) of the LCA according to standard procedures (10), described in the Supplemental Methods. Following 24-hour reperfusion, hearts of surviving mice were infiltrated with Evans blue and removed for histological analysis.

**TAC model.** TAC was carried out as described (15) in male mice 9 to 16 weeks of age. In brief, the aorta was exposed following induction of anesthesia and a ligature of 7-0 silk suture was placed around the aortic arch and an apposed segment of 27-gauge needle, between the right innominate and left carotid arteries. The ligature was tightened, and the needle was removed immediately. The resulting reduction in the cross-sectional area of the arch was approximately 90%.

**Echocardiography.** Mice were chemically restrained with isoflurane, and transthoracic echocardiography was performed using a Vevo 770 High-Resolution Imaging System equipped with an RMV-707B 30-MHz probe (VisualSonics). Standard M-mode sampling was used through the LV short axis at the midpapillary level. Ejection fraction, fractional shortening, and other parameters were determined using the system's software. LV wall thickening was calculated as follows: (end systolic wall thickness — end diastolic wall thickness)/end diastolic wall thickness × 100%.

**Statistics.** Data are presented as means (± SEM) or as ratios with group differences tested using standard parametric methods (i.e. Stu-

dent's *t* test, 2-tailed) or 2 × 2 contingency tables with Fisher's exact test as appropriate. In all settings, a *P* value of less than 0.05 was assumed to indicate a statistically significant difference in the parameter being compared.

**Study approval.** All experimental procedures were approved by the Institutional Animal Care and Use Committee of Case Western Reserve University School of Medicine and were conducted in accordance with the NIH *Guide for the Care and Use of Laboratory Animals* (National Academies Press, 2011). All potentially painful procedures were conducted under a surgical plane of anesthesia (typically ketamine/xylazine 100/10 mg/kg), and mice were killed by methods compliant with the American Veterinary Medical Association (AMVA) *Guidelines for Euthanasia of Animals* (2013 edition).

## Author contributions

RZ, DTH, and JSS designed the studies; RZ conducted the experiments; RZ, DTH, and JDR analyzed the data; all authors participated in writing the manuscript.

## Acknowledgments

This work was supported in part by NIH grants 5P01HL075443 and 1R01HL126900-01A1. The mouse strains employed were provided by T.M. Townes (University of Alabama, Birmingham, Alabama, USA).

Address correspondence to: Jonathan S. Stamler, Institute for Transformative Molecular Medicine, Case Western Reserve University, 10900 Euclid Avenue, Location Code 7294, Cleveland, Ohio 44106, USA. Phone: 216.368.5725; E-mail: jonathan.stamler@case.edu.

- Ross JM, Fairchild HM, Weldy J, Guyton AC. Autoregulation of blood flow by oxygen lack. *Am J Physiol.* 1962;202:21-24.
- Stamler JS, et al. Blood flow regulation by S-nitrosohemoglobin in the physiological oxygen gradient. *Science.* 1997;276(5321):2034-2037.
- Jia L, Bonaventura C, Bonaventura J, Stamler JS. S-nitrosohaemoglobin: a dynamic activity of blood involved in vascular control. *Nature.* 1996;380(6571):221-226.
- Singel DJ, Stamler JS. Chemical physiology of blood flow regulation by red blood cells: the role of nitric oxide and S-nitrosohemoglobin. *Annu Rev Physiol.* 2005;67:99-145.
- Zhang R, et al. Hemoglobin βCys93 is essential for cardiovascular function and integrated response to hypoxia. *Proc Natl Acad Sci U S A.* 2015;112(20):6425-6430.
- Isbell TS, et al. SNO-hemoglobin is not essential for red blood cell-dependent hypoxic vasodilation. *Nat Med.* 2008;14(7):773-777.
- Nagai K, Perutz MF, Poyart C. Oxygen binding properties of human mutant hemoglobins synthesized in *Escherichia coli*. *Proc Natl Acad Sci U S A.* 1985;82(21):7252-7255.
- Parodi O, et al. Myocardial blood flow distribution in patients with ischemic heart disease or dilated cardiomyopathy undergoing heart transplantation. *Circulation.* 1993;88(2):509-522.
- Briceno N, Schuster A, Lumley M, Perera D. Ischaemic cardiomyopathy: pathophysiology, assessment and the role of revascularisation. *Heart.* 2016;102(5):397-406.
- Michael LH, et al. Myocardial ischemia and reperfusion: a murine model. *Am J Physiol.* 1995;269(6 Pt 2):H2147-H2154.
- Rockman HA, et al. Segregation of atrial-specific and inducible expression of an atrial natriuretic factor transgene in an in vivo murine model of cardiac hypertrophy. *Proc Natl Acad Sci U S A.* 1991;88(18):8277-8281.
- Gaston B, et al. Essential role of hemoglobin beta-93-cysteine in posthypoxia facilitation of breathing in conscious mice. *J Appl Physiol.* 2014;116(10):1290-1299.
- Sano M, et al. p53-induced inhibition of Hif-1 causes cardiac dysfunction during pressure overload. *Nature.* 2007;446(7134):444-448.
- Yang J, Gonon AT, Sjöquist PO, Lundberg JO, Pernow J. Arginase regulates red blood cell nitric oxide synthase and export of cardioprotective nitric oxide bioactivity. *Proc Natl Acad Sci U S A.* 2013;110(37):15049-15054.
- Hu P, Zhang D, Swenson L, Chakrabarti G, Abel ED, Litwin SE. Minimally invasive aortic banding in mice: effects of altered cardiomyocyte insulin signaling during pressure overload. *Am J Physiol Heart Circ Physiol.* 2003;285(3):H1261-H1269.